

## Letter to the Editor

# Refined Genetic Mapping of X-Linked Thoracoabdominal Syndrome

### To the Editor:

We have previously described thoracoabdominal syndrome [TAS; Carmi et al., 1990] and mapped the gene to Xq25-26 [Parvari et al., 1994]. After using new polymorphic markers made available since the previous publication, and detecting a reading error of one allele of a CA repeat at the DXS102 locus in one individual in the affected family, we refined the localization of TAS to Xq27, with no recombination observed with DXS1232 and DXS984.

Microsatellite polymorphic loci available for the Xq25-q27 interval were tested on the 27 relatives of the TAS family. Linkage between the TAS locus and marker loci was calculated using ILINK and MLINK from the Linkage program v5.1 [Lathrop et al., 1984]. The TAS locus was modeled as an X-linked dominant locus with full penetrance. The pairwise linkage data for the informative markers on Xq25-q27 are presented in Table I. The highest lod scores of 5.49 and 6.36 were obtained for DXS1232 and DXS984, respectively, at  $\theta = 0$ . The first neighboring loci showing one cross-over were F9 on the centromeric side and DXS1205 on the telomeric side (the latter showing a different recombination event). The estimated genetic distance between these flanking loci is about 5 cM and the physical distance is 2.5 Mb ( $\pm 100$  Kb) [Little et al., 1991; Zucchi et al., manuscript in preparation]. In addition to the loci presented in Table I the following loci were tested and found to be noninformative: DXS1062, DXS1211, DXS998, DXS1215, DXS1193 [Gyapay et al., 1994], and AFM144yd2 (Genethon). Additional uninformative restriction fragment length polymorphism (RFLP) loci tested included: DXS105 (probe cX55, DNA digested with EcoR I, Taq I, and Hind III) [Hofker et al., 1985]; DXS 152 (probe cX33.2, DNA digested with ApaI I) [Hofker et al., 1985]; and DXS 119 (probe 780, DNA digested with Bcl I) [Hofker et al., 1985]. The results presented here thus narrow down the region for the search for the TAS gene to an interval of about 2.5 Mb. SOX3, a gene related to SRY and expressed in mouse embry-

onal tissues, has been mapped to this interval [Stevanovich et al., 1993] and is a candidate gene. Genes for several other disorders have been localized in the vicinity of the TAS interval. Among them are premature ovarian failure with a deletion of Xq26.1-28 presenting with no other abnormalities [Tharapel et al., 1993]; split hand split foot in Xq26.1 [Faiyaz ul Haque et al., 1992]; Simpson-Golabi-Behmel syndrome in Xq25-27 [Xuan et al., 1994]; and a heterotaxy gene has also been placed somewhere in Xq24-q27 [Casey et al., 1993]. This latter condition of faulty body lateralization is of special interest in our context, since midline formation and body placement determination are developmentally related blastogenetic events. It is conceivable that the region Xq24-q27 contains a cluster of developmental genes that operate in a concerted fashion.

### ACKNOWLEDGMENTS

We are grateful to Dr. David Schlessinger for the kind gift of probes cX55, cX33.2, and 780. This research is supported by the Basic Research Foundation administered by the Israeli Academy of Sciences and Humanities and the Ministry of Science and Technology.

### REFERENCES

- Carmi R, Barbash A, Mares AJ (1990): The thoracoabdominal syndrome (TAS): A new X-linked dominant disorder. *Am J Med Genet* 36:109–114.
- Casey B, Devoto M, Jones KL, Ballabio A (1993): Mapping a gene for familial situs abnormalities to human chromosome Xq24-q27.1. *Nature Genet* 5:403–407.
- Faiyaz ul Haque M, Uhlhaas S, Knapp M, Schuler H, Friedl W, Ahmad M, Propping P (1992): X-chromosomal split-hand/split-foot anomaly maps to Xq26. *Am J Hum Genet* 51(Suppl):A189.
- Gedeon AK, Holman K, Richards RI, Mulley JC (1992): Characterization of new PCR based markers for mapping and diagnosis: AC dinucleotide repeat markers at the DXS237 (GMGX9) and DXS102 (cX38.1) loci. *Am J Med Genet* 43:255–260.
- Graham JB, Kunkel GR, Egilmez NK, Wallmark A, Fowlkes DM, Lord ST (1991): The varying frequencies of five DNA polymorphisms of X-linked coagulant factor IX in eight ethnic groups. *Am J Hum Genet* 49:537–544.
- Gyapay G, Morissette J, Vignal A, Dib C, Fizames C, Millasseau P, Marc S, Bernardi G, Lathrop M, Weissenbach J (1994): The 1993–1994 Genethon human genetic linkage map. *Nature Genet* 7:246–339.
- Hearne CM, Todd JA (1991): Tetranucleotide repeat polymorphism at the HPRT locus. *Nucleic Acids Res* 19:5450.
- Hofker MH, van Ommen CJB, Bergen A, Skraastad MI, Bakker E, Ropers HH, Pearson PL (1985): Strategy for efficient X-probe isolation via cosmid libraries; Detection of FFLPs on the X-chromosome. *Cytogenet Cell Genet* 40:652–653.

Received for publication November 15, 1994; revision received August 18, 1995.

Address reprint requests to Rivka Carmi, MD, Clinical Genetics Unit, Soroka Medical Center, P.O. Box 151, Beer-Sheva 84101, Israel.

TABLE I. Two-Point Linkage Analysis of TAS to Xq25-27 Markers

TABLE 1. Two Point Linkage Analysis of the 10 Locus 27 Markers										
Recombination fraction										
	0.00	0.01	0.05	0.10	0.20	0.30	0.40	$\theta$ max	Z max	References#
DXS1047	$-\infty$	5.33	5.54	5.21	4.14	2.79	1.32	0.04	5.57	Gyapay et al. [1994]
HPRT	$-\infty$	5.22	5.48	5.21	4.26	3.03	1.58	0.04	5.50	Hearne and Todd [1991]
DXS994	$-\infty$	4.82	5.11	4.87	4.00	2.85	1.50	0.04	5.12	Weissenbach et al. [1992]
DXS1192	$-\infty$	4.11	4.41	4.20	3.38	2.30	1.10	0.04	4.42	Gyapay et al. [1994]
DXS102	$-\infty$	4.81	5.09	4.83	3.92	2.73	1.36	0.04	5.10	Gedeon et al. [1992]
F9	$-\infty$	3.38	3.79	3.69	3.11	2.27	1.23	0.06	3.79	Graham et al. [1991] <sup>a</sup>
DXS1232	5.49	5.41	5.10	4.66	3.72	2.64	1.41	0.00	5.49	Gyapay et al. [1994]
DXS984	6.36	6.25	5.83	5.27	4.06	2.71	1.27	0.00	6.36	Weissenbach et al. [1992]
DXS1205	$-\infty$	3.75	4.06	3.85	3.10	2.20	1.19	0.04	4.06	Gyapay et al. [1994]
DXS1227	$-\infty$	4.57	4.84	4.58	3.66	2.45	1.04	0.04	4.86	Gyapay et al. [1994]

<sup>a</sup> Taq I polymorphism.

Lathrop GM, Lalouel JM, Julier C, Ott J (1984): Strategies for multi-locus linkage analysis in humans. *Proc Natl Acad Sci USA* 81:3443–3446.

Little RD, Pilia G, Johnson S, D'Urso M, Schlessinger D (1991): Yeast artificial chromosomes spanning 8 megabases and 10–15 centimorgans of human cytogenetic band Xq26. *Proc Natl Acad Sci USA* 89:177–181.

Parvari R, Weinstein Y, Ehrlich S, Steinitz M, Carmi R (1994): Linkage localization of the thoraco-abdominal syndrome (TAS) gene to Xq25-26. *Am J Med Genet* 49:431–434.

Stevanovich M, Lovell-Badge R, Collignon J, Goodfellow PN (1993): SOX3 is an X-linked gene related to SRY. *Hum Mol Genet* 12:2013–2018.

Tharapel AT, Anderson KP, Simpson JL, Martens PR, Wilroy RS Jr, Llerena JC Jr, Schwartz CE (1993): Deletion (X) (q26.1→q28) in a proband and her mother: Molecular characterization and phenotypic-karyotypic deductions. *Am J Hum Genet* 52:463–471.

Weissenbach J, Gyapay G, Dib C, Vignal A, Morissette J, Millasseau P, Baysseix G, Lathrop M (1992): A second generation linkage map of the human genome. *Nature* 359:794–801.

Xuan JY, Besner A, Ireland M, Hughes-Benzie RM, MacKenzie AE (1994): Mapping of Simpson-Golabi-Behmel syndrome to Xq25-27. *Hum Mol Genet* 3:133–137.

**R. Parvari**  
**R. Carmi**  
Clinical Genetics Unit

**J. Weissenbach**  
Genethon  
Evry, France

**G. Pilia**  
**S. Mumm**  
Center for Genetics in Medicine  
Washington University School of Medicine  
St. Louis, Missouri

**Y. Weinstein**  
Department of Microbiology and Immunology  
Soroka Medical Center  
Ben-Gurion University of the Negev  
Beer-Sheva, Israel